

CFT description of the Fully Frustrated XY model and phase diagram analysis

Gerardo Cristofano¹, Vincenzo Marotta¹, Petter Minnhagen², Adele Naddeo³, Giuliano Niccoli⁴.

Abstract

Following a suggestion given in [1], we show how the $U(1) \otimes Z_2$ symmetry of the fully frustrated XY (FFXY) model on a square lattice can be accounted for in the framework of the m -reduction procedure developed for a Quantum Hall system at “paired states” fillings $\nu = 1$ [2, 3]. The resulting twisted conformal field theory (CFT) with central charge $c = 2$ is shown to well describe the physical properties of the FFXY model. In particular the whole phase diagram is recovered by analyzing the flow from the Z_2 degenerate vacuum of the $c = 2$ CFT to the infrared fixed point unique vacuum of the $c = \frac{3}{2}$ CFT. The last theory is known to successfully describe the critical behavior of the system at the overlap temperature for the Ising and vortex-unbinding transitions.

Keywords: Twisted CFT, Fully Frustrated XY model

PACS: 11.25.Hf, 02.20.Sv, 03.65.Fd

¹Dipartimento di Scienze Fisiche, Università di Napoli “Federico II”
and INFN, Sezione di Napoli, Via Cintia, Compl. universitario M. Sant’Angelo, 80126 Napoli, Italy

²Department of Theoretical Physics, Umea University, 901 87 Umea, Sweden

³Dipartimento di Fisica “E. R. Caianiello”, Università degli Studi di Salerno
and CNISM, Unità di Ricerca di Salerno, Via Salvador Allende, 84081 Baronissi (SA), Italy

⁴Laboratoire de physique, Ecole Normale Supérieure de Lyon, 46 allée d’Italie, 69364 Lyon cedex 07, France.

1 Introduction

A few years ago Foda [1] proposed that the continuum limit of a FFX model is equivalent, at the overlap temperature, to a free massless field theory of one boson and one Majorana fermion, that is a superconformal field theory with central charge $c = 3/2$. Such a model shares both discrete and continuous symmetries, which describe well a Josephson junction array in a transverse magnetic field, provided that the external magnetic flux threading each cell of the array is $\frac{1}{2}\Phi_0$, where $\Phi_0 = \frac{hc}{2e}$ is the superconducting flux quantum [4, 5].

The XY model on a square lattice in the presence of an external magnetic field transversal to the lattice plane is described by the action:

$$H = -\frac{J}{kT} \sum_{\langle ij \rangle} \cos(\varphi_i - \varphi_j - A_{ij}), \quad (1)$$

where $\{\varphi\}$ are the phase variables defined on the sites, the sum is over nearest neighbors, $J > 0$ is the coupling constant and $A_{ij} = \frac{2e}{\hbar c} \int_i^j A \cdot dl$ is the line integral along the bond between adjacent sites i and j . We consider the case where the bond variables A_{ij} are fixed, uniformly quenched, out of equilibrium with the site variables and satisfy the condition $\sum_p A_{ij} = 2\pi f$; here the sum is over each set of bonds of an elementary plaquette and f is the strength of frustration. We assume that the local magnetic field in Eq. (1) is equal to the uniform applied field; such an approximation is more valid the smaller is the sample size L compared with the transverse penetration depth $\lambda_\perp$ [6]. In the case of full frustration, i.e. $f = \frac{1}{2}$, of interest to us here, such a model has a continuous $U(1)$ symmetry associated with the rotation of spins and an extra discrete Z_2 symmetry, as it has been shown analysing the degeneracy of the ground state [7, 8]. Choosing the Landau gauge, such that the vector potential vanishes on all horizontal bonds and on alternating vertical bonds, we get a lattice where each plaquette displays one antiferromagnetic and three ferromagnetic bonds. Such a choice corresponds to switching the sign of the interaction term in Eq. (1) and is closely related to the presence of two ground states with opposite chiralities, the first one invariant under shifts by two lattice spacings and the second one invariant under shifts by one lattice spacing.

In the presence of such a degeneracy we can use the m -reduction technique [3] which has been successfully applied to a quantum Hall fluid in [2, 3, 9, 10, 11] and to a fully frustrated Josephson junction ladder with Mobius boundary conditions in [12]. In Section 4 we will see how the discrete version of the m -reduction procedure, successfully used to build a CFT with a Z_m -twist, well describes such a situation.

The Z_2 symmetry of the FFX model is broken at low temperature and will be restored beyond a certain temperature after the formation of domain walls separating islands of opposite chirality. The Ising transition overlaps to a vortex-unbinding transition [13], which is associated with the $U(1)$ symmetry [8, 14].

The action (1) can be cast into a form where both the $U(1)$ and Z_2 symmetries are manifest, through the Villain approximation [15]. In this way the spin-wave and the

vortex contributions can be separated. Furthermore, by integrating out the spin waves, the resulting vortex contribution can be rewritten as a fractionally charged Coulomb gas (CG) defined on the dual lattice [8, 16]:

$$H = -\frac{J}{kT} \sum_{r,r'} (m(r) + f) G(r, r') (m(r') + f). \quad (2)$$

Here we have $\lim_{|r-r'|\rightarrow\infty} G(r, r') = \log|r-r'| + \frac{1}{2}\pi$ and the neutrality condition $\sum_r (m(r) + f) = \sum_r n(r) = 0$ must be satisfied. It is now evident that the ground state for $f = \frac{1}{2}$ consists of an alternating lattice of logarithmically interacting $\pm\frac{1}{2}$ charges and is doubly degenerate. Such a model exhibits two possible phase transitions, an Ising and a vortex-unbinding one, and their relative order has been deeply studied in the literature [7].

It is just the doubly degenerate checkerboard pattern of vortices in the ground state that gives rise to the Ising-like Z_2 discrete symmetry in addition to the Kosterlitz-Thouless (KT) continuous $U(1)$ one, associated with the uniform rotation of all phase angles [13]. The issue whether there are two distinct phase transitions, $T_V > T_{ISING}$ or $T_V < T_{ISING}$ with T_V and T_{ISING} marking respectively the breaking of $U(1)$ and of Z_2 symmetry [17], or a single transition with the simultaneous breaking of both symmetries [18] has been widely investigated. Indeed an exact mapping of the classical 2D FFX model onto an alternating 19-vertex model has been established [19], whose equivalence with the restricted solid-on-solid (RSOS) model is well known [20]. In turn the RSOS model can be viewed as a discretized time generalization of the spin-1 one dimensional quantum chain, as shown in detail in Ref. [21] by means of the transfer matrix formalism. The phase diagram of such a system has been widely investigated in a simplified case, i.e. the XXZ quantum chain with single-ion-type anisotropy, by means of analytical and numerical techniques [21, 22, 23]. It shows both $U(1)$ and Z_2 symmetry breaking in the space $D - J_z$ (where the parameter D is the uniaxial single-ion anisotropy) [21, 24, 25]. Depending on the values of the parameters different phases of the system emerge: the Haldane phase, the large- D phase, two XY phases (XY1 and XY2) with different Z_2 symmetry, the ferromagnetic phase and the Neel phase. Then various types of phase transitions take place. In particular all the three possibilities above quoted can be recognized: a gapful-gapless KT transition between the Haldane phase and the XY1 one followed by a transition between the two different XY phases, XY1 and XY2; an Ising transition between the Haldane phase and Neel phase and after that a transition Neel-XY2 at $J_z = 0$ for large negative D ; an XY and Ising transition lines which merge at a tricritical point within $c = \frac{3}{2}$ CFT universality class. All that reveals the richness of the phase structure of a class of frustrated systems with the same ground state but the controversy on the phase diagram structure of the FFX model remains still unsolved.

Recently Korshunov [26] argued that a new transition could take place in the 2D FFX model, well below the bulk transition, due to the unbinding of kink-antikink pairs on the domain walls associated with Z_2 symmetry. Such a transition could lead to a decoupling of phase coherence across domain boundaries, so producing two distinct bulk transitions

with $T_V < T_{ISING}$ and then a coupled XY-Ising model [27, 28]. Direct numerical evidence supporting the existence of such a transition at a temperature T_W below the bulk ones has by now been provided [29]. Also Teitel and Jayaprakash [14] have considered the case $T_V < T_{ISING}$ in terms of a dual Coulomb gas model. They argued that the helicity modulus Y would decrease from its $T = 0$ value by means of fluctuations producing dipole moments. So, if Ising domains of size ξ_{ISING} carried a total dipole moment proportional to their size, they would make $Y = 0$ in a continuous way, due to the divergence of ξ_{ISING} as $T \rightarrow T_{ISING}$. At $T = 0$ it has been found though that the only Ising-like domains are those which carry no dipole moment: they cannot produce a reduction in Y . But at $T > T_W$, according to Ref. [26], kink-antikink excitations on the boundaries of the domain walls unbind, so the Ising domains would acquire large dipole moments without cost in free energy. Such a conclusion [29] supports strongly the existence of two separate transitions with $T_V < T_{ISING}$. This scenario has been recently confirmed also by numerical simulations on very large lattices and by means of a careful finite-size scaling analysis in Refs. [30]. The critical behaviour at the overlap temperature is well described by a conformal field theory with central charge $c = 3/2$ [1]. The interesting scenario just outlined is the result of an interplay between the Ising and gaussian sectors of the model: they are linked through the fractional vortices which reside at the corners of the domain walls in the off-critical theory at the microscopic level [7]. More recent Monte Carlo simulations of 2D XY-type models [31] appear to confirm the above scenario [32]. Indeed the sequence of phase transitions on going from higher to lower temperatures is: first Ising, then KT and next kink-antikink unbinding. This picture will then remain (with Ising and KT transition very close to each other) until a particular value of the parameters where all the three lines merge at the same time. After that the transition is first order and the critical point is characterized by a central charge $c = 2$ [32].

The FFXY model has an interesting experimental realization as a Josephson junctions array (JJA) in a transverse magnetic field with half a magnetic flux quantum per cell [33, 34]. Measurements have been performed on the electrical properties of such a system, in particular on the resistance as a function of the temperature and the external magnetic field which is closely related to the helicity modulus and then to the vortex-unbinding transition [33]. The fact that the vortex transition is shifted downwards in temperature [33] is in agreement with the results found in Ref. [35], which refer to an XY model with a modified potential of the form:

$$V(\varphi) = \cos \varphi + \alpha \cos 2\varphi. \quad (3)$$

That tentatively leads to the conclusion that the arrays of Ref. [33] display all the properties of the FFXY model with higher harmonics contributions to the potential [1].

In this paper we construct a CFT for such systems which extends the results of Ref. [1], so recovering the whole phase diagram quoted in Refs. [27, 28, 36, 37], then we close with a brief account on the relation between the FFXY model and the physics of a Josephson junction array in the classical approximation [38].

The paper is organized as follows.

In Section 2 we discuss in a qualitative way the phase diagram of the FFXY model describing some models which have the same $U(1) \otimes Z_2$ degenerate ground state and are believed to be in the same universality class.

In Section 3 we recall some aspects of the m -reduction procedure, in particular we show how the $m = 2$, $p = 0$ case well accounts for the $U(1) \otimes Z_2$ symmetry of the FFXY model. In such a framework we give the whole primary fields content of the theory on the plane.

In Section 4 we give a discrete version of the m -reduction procedure and then, starting from our CFT results, we show how to recover the whole FFXY phase diagram introduced in Section 2.

In Section 5 some comments and outlooks are given.

2 Phase diagram of the FFXY model

In the literature the phase transitions of the FFXY model have been studied analyzing other models which have the same $U(1) \otimes Z_2$ degenerate ground state and are believed to be in the same universality class. In particular, by using symmetry arguments[7, 18] or a Hubbard-Stratonovich transformation[27], the FFXY model can be reformulated in terms of a system of two coupled XY models with a symmetry breaking term:

$$H = A \left[\sum_{i=1,2} \sum_{\langle r, r' \rangle} \cos \left(\varphi^{(i)}(r) - \varphi^{(i)}(r') \right) \right] + h \sum_r \cos 2 \left(\varphi^{(1)}(r) - \varphi^{(2)}(r) \right). \quad (4)$$

Such a system has been studied in detail by Granato and Kosterlitz [39]. The limit $h \rightarrow 0$ corresponds to a full decoupling of the fields $\varphi^{(i)}$, $i = 1, 2$, so giving rise to a CFT with central charge $c = 2$ which describes two independent classical XY models (in the continuum limit). In the $h \rightarrow \infty$ limit, the two phases $\varphi^{(i)}$, $i = 1, 2$ are locked [39]:

$$\varphi^{(1)}(r) - \varphi^{(2)}(r) = \pi j, \quad j = 1, 2; \quad (5)$$

as a consequence the model gains a symmetry $U(1) \otimes Z_2$ and its Hamiltonian renormalizes towards a model described by:

$$H = H(h \rightarrow \infty) = A \sum_{\langle r, r' \rangle} (1 + s_r s_{r'}) \cos \left(\varphi^{(1)}(r) - \varphi^{(1)}(r') \right) \quad (6)$$

where $\varphi^{(1)}(r)$ and $s_r = \cos \pi j = \pm 1$ are planar and Ising spins respectively. In this way a model is obtained which is consistent with the required symmetry, the XY-Ising one and whose Hamiltonian has the general form [27, 36]:

$$H_{XY-I} = \sum_{\langle r, r' \rangle} \left[A(1 + s_r s_{r'}) \cos \left(\varphi^{(1)}(r) - \varphi^{(1)}(r') \right) + C s_r s_{r'} \right], \quad (7)$$

which is believed to be in the universality class of the classical FFXY model. However such a model does not contain fractional vortices at the corners of the domain walls [7], which are believed to give rise to an interesting interplay between the continuous and discrete symmetries [1]. It is just this complex interplay which accounts for the unusual and not yet fully understood critical behaviour of the FFXY model on the square lattice [1, 27, 28, 36, 37], that will be discussed in the following.

The phase diagram of the model [27, 28, 36, 37] with Hamiltonian (7), as depicted in Fig. 1, is built up with three branches which meet at a multi-critical point P . The branch PT corresponds to single transitions with simultaneous loss of XY and Ising order while the other two describe separate Kosterlitz-Thouless and Ising transitions. The critical line PT becomes a first order one at the tricritical point P ; it seems to be non-universal [27, 40] as the values of the critical exponents ($\nu = 0.79$, $\eta = 0.40$) estimated by finite-size scaling of large systems are inconsistent with pure Ising critical behavior ($\nu = 1$, $\eta = \frac{1}{4}$). Also the numerical estimate for the central charge c ($c \sim 1.60$) [28] is higher than the expected value ($c = 3/2$) for a critical behavior given by the superposition of critical Ising ($c = 1/2$) and gaussian ($c = 1$) models [1]. Indeed the central charge seems to vary continuously from $c \approx 1.5$ near P to $c \approx 2$ at T [28]. All such results would suggest the existence of a parameter changing along the critical PT line that does not affect the symmetry [37]. The system seems to be not conformally invariant [27], so it would be natural to consider all possible integrable perturbations of a superposition of critical Ising and XY models as a starting point to study the vicinity of the point P [37]. Furthermore different numerical techniques have led to different results suggesting either single [18] or double transitions [17] along the critical PT line. How to reconcile such conflicting results? A possible solution is suggested in Ref. [41] where the Coulomb gas Hamiltonian of Eq. (2) gets modified through an additional antiferromagnetic coupling between nearest-neighbour vortices:

$$H = -\frac{J}{kT} \sum_{r, r'} n_r G(r, r') n_{r'} + \bar{J} \sum_{\langle r, r' \rangle} n_r n_{r'}. \quad (8)$$

Such a model has a phase diagram with a structure close to the XY-Ising model one of Fig. 1 with the occurrence of double or single transitions on the PT line depending on this additional coupling; for $\bar{J} \neq 0$ the two transitions separate with the vortex-unbinding one occurring at a lower temperature [41]. A similar effect has been observed theoretically in Ref. [35], where a XY model with various modifications of the cosine-type potential




Figure 1: The phase diagram for the XY-Ising model [27, 36].

is studied, and experimentally in Ref. [33]: it can be ascribed to the presence of higher harmonics contributions to the potential. Such a finding is a crucial one, as we will show in Section 4.

Another route to the solution of all these issues is the m -reduction technique [2, 3] which describes well the two fixed points with $c = 3/2$ and $c = 2$ and accounts for non trivial boundary conditions for the square lattice. Such a realization of the FFXY model on closed geometries could be relevant for the description of JJAs with non trivial topologies, which are believed to provide a physical implementation of an ideal quantum computer [42] because of the topological ground state degeneracy “protected” from external perturbations [43]. We introduce the m -reduction in the next Section.

3 A review of the m -reduction procedure

In this Section we recall those aspects of the twisted model (TM) which are relevant for the FFXY model. We focus in particular on the m -reduction procedure for the special $m = 2$ case (see Ref. [2] for the general case), since we are interested in a system with $U(1) \otimes Z_2$ symmetry. We showed in Refs. [2, 3] that such a theory describes well a system consisting of two parallel layers of 2D electron gas in a strong perpendicular magnetic field, with filling factor $\nu^{(a)} = \frac{1}{2p+2}$ for each of the two $a = 1, 2$ layers and total filling $\nu = \nu^{(1)} + \nu^{(2)} = \frac{1}{p+1}$. Regarding the integer p , characterizing the flux attached to the particles, we choose the “bosonic” value $p = 0$, since it enables us to describe the highly correlated system of vortices with flux quanta $\frac{hc}{2e}$.

Let us start from the bosonic “filling” $\nu = \frac{1}{2}$, described by a CFT with $c = 1$ in terms of a scalar chiral field Q compactified on a circle with radius $R^2 = 1/\nu = 2$. It is explicitly given by:

$$Q(z) = q - i p \ln z + i \sum_{n \neq 0} \frac{a_n}{n} z^{-n}, \quad (9)$$

with a_n , q and p satisfying the commutation relations $[a_n, a_{n'}] = n \delta_{n,n'}$ and $[q, p] = i$.

From such a CFT (mother theory), using the m -reduction procedure, which consists in considering the subalgebra generated only by the modes in Eq. (9) which are a multiple of an integer m , we get a $c = m$ orbifold CFT (daughter theory, i.e. the TM) which, for $m = 2$, in the next Section will be shown to describe the whole phase diagram of the FFXY model. Then the fields in the mother CFT can be organized into components which have well defined transformation properties under the discrete Z_m (twist) group, which is a symmetry of the TM. By using the mapping $z \rightarrow z^{1/m}$ and by making the identifications $a_{nm+l} \rightarrow \sqrt{m} a_{n+l/m}$, $q \rightarrow \frac{1}{\sqrt{m}} q$, the $c = m$ CFT (daughter theory) is obtained.

Its primary fields content, for the special $m = 2$ case, can be expressed in terms of a Z_2 -invariant scalar field $X(z)$, given by

$$X(z) = \frac{1}{2} (Q(z) + Q(-z)), \quad (10)$$

describing the electrically “charged” sector of the new theory, and a twisted field

$$\phi(z) = \frac{1}{2} (Q(z) - Q(-z)), \quad (11)$$

which satisfies the twisted boundary conditions $\phi(e^{i\pi}z) = -\phi(z)$ and describes the “neutral” sector [2].

The TM primary fields are composite vertex operators $V(z) = U_X(z) \psi(z)$ where

$$U_X(z) = \frac{1}{\sqrt{z}} : e^{i\alpha X(z)} : , \quad \alpha^2 = 2 \quad (12)$$

is the vertex of the “charged” sector and

$$\psi(z) = \frac{1}{\sqrt{z}} : e^{i\alpha\phi(z)} : \quad (13)$$

is the “neutral” one. We point out that in the neutral sector it is useful to introduce the two chiral operators:

$$\psi(z) = \frac{1}{2\sqrt{z}} (: e^{i\alpha\phi(z)} : + : e^{i\alpha\phi(-z)} :), \quad \bar{\psi}(z) = \frac{1}{2\sqrt{z}} (: e^{i\alpha\phi(z)} : - : e^{i\alpha\phi(-z)} :); \quad (14)$$

the first one does not change the boundary conditions while the second one does. In a fermionized version of the theory they correspond to two $c = 1/2$ Majorana fermions with Ramond and Neveu-Schwartz boundary conditions [2, 3]. Furthermore in the TM they appear to be not completely equivalent.

In fact the whole TM theory decomposes into a tensor product of two CFTs, a twisted invariant one with $c = 3/2$ (the Moore-Read (MR) theory with symmetry $U(1) \otimes Z_2$) and the remaining $c = 1/2$ one realized by a Majorana fermion in the twisted sector. Such a factorization can be unambiguously pointed out on the torus topology where the partition function of the $c = 2$ theory, i.e. the TM, can be written as⁵:

$$Z(w_c|\tau) = Z^{MR}(w_c|\tau) Z_{\bar{T}}(\tau). \quad (15)$$

Here $Z^{MR}(w_c|\tau)$ stands for the MR partition function and $Z_{\bar{T}}(\tau)$ is the partition function describing the Ising degrees of freedom which are Z_2 antisymmetric (i.e. $\bar{\psi}(z)$ of Eq. (14) in the plane geometry).

⁵See Appendix for the explicit expression and Ref. [3] for its derivation.

Let us point out that the energy-momentum tensor of the Ramond part of the neutral sector develops a cosine term:

$$T_\psi(z) = -\frac{1}{4}(\partial\phi)^2 - \frac{1}{16z^2}\cos\left(2\sqrt{2}\phi\right), \quad (16)$$

a clear signature of a tunneling phenomenon which selects out a new stable vacuum, the $c = 3/2$ one. We identify such a Z_2 invariant theory with the one describing the FFX model conjectured in Ref. [1]. Here we only point out that the critical behavior of the $c = 3/2$ theory will be a superposition of the critical behavior of its components with $c = 1$ and $c = 1/2$ respectively. The gaussian component will lead to a gaussian critical line that ends at a vortex-unbinding transition point, while the Ising component will lead to an Ising critical point. The conformal field theory, considered in this paper, is not meant to predict the position of the Ising point with respect to the gaussian critical line, it only tells us that there exists an Ising transition overlapping somewhere with a gaussian critical line. In the following we exhibit all the relevant physical consequences of our theory.

4 TM description of the FFX phase diagram

In this Section we will derive the phase diagram of the FFX model⁶ in terms of the RG flow which originates from perturbing our TM model.

As we pointed out in Section 2, the limit $h \rightarrow 0$ in the Hamiltonian of Eq. (4) corresponds to a full decoupling of the fields $\varphi^{(i)}$, $i = 1, 2$. In the continuum limit that gives rise to a CFT with two scalar Fubini boson fields $\varphi^{(i)}$ and with central charge $c = 2$. Let us stress that a good candidate to describe the FFX model at criticality around the point T of the phase diagram is a CFT, with $c = 2$, which accounts for the full spectrum of excitations of the model. So it is worth recalling briefly the structure of such excitations for the FFX model. A domain wall is a topological excitation of the double degenerate ground state and it can be defined as a line of links, each one separating two plaquettes with the same chirality. So, through a domain wall the alternating structure (the checkerboard pattern) of the ground state is lost. Kinks and antikinks are excitations which live on the domain walls and are described by fractional vortices with $+1/2$ and $-1/2$ topological charge. Such particle-like excitations belong also to the spectrum of the fully frustrated ladders of Josephson junctions and in [12] we have shown how the TM model exactly describes them. In particular these excitations can be generated from the ground state by closing the ladder, by imposing the coincidence of opposite sides. In fact, in this closed geometry, for an even number of plaquettes we obtain the ground state while for the odd case this point-like excitation is generated on the boundary. In the TM model they are described by different boundary conditions, untwisted and twisted ones, respectively.

⁶As it has been described in Section 2 and inferred by Migdal-Kadanoff renormalization [36], Monte Carlo simulations [27] and Monte Carlo transfer matrix calculations [28].

Similar considerations apply to the FFX array; in particular the central role played by the closed geometry and by the boundary conditions in the description of the excitation spectrum is clear already at the level of the lattice model. By imposing the coincidence between opposite sides of the square lattice, we obtain a closed geometry, which is the discretized analogue of a torus. Now for the ground state two topologically inequivalent circumstances arise, one for even and the other one for odd number of plaquettes. In the even case the end plaquettes on the opposite sides of the lattice have opposite chirality, while in the odd case they have the same chirality. So the ground state on the square lattice maps into the ground state for the even case while it generates two straight domain walls along the two cycles of the torus for the odd case. Such a behaviour has to be taken into account by opportune boundary conditions on the field $\varphi^{(i)}$ at the borders of the finite lattice. These non trivial boundary conditions naturally arise when we implement the m -reduction procedure in the discrete case. To this aim, let $(-L/2, 0)$, $(L/2, 0)$, $(L/2, L)$, $(-L/2, L)$ be the corners of the square lattice $\mathcal{L}$ and assume that the fields $\varphi^{(i)}$ satisfy the following boundary conditions:

$$\varphi^{(1)}(r) = \varphi^{(2)}(r) \quad \text{for } r \in \mathcal{L} \cap \mathbf{x}, \quad (17)$$

where $\mathbf{x}$ is the x axis. The above boundary conditions allow us to consider the two fields $\varphi^{(1)}$ and $\varphi^{(2)}$ on the square lattice $\mathcal{L}$ as the folding of a single field $\mathcal{Q}$, defined on the lattice $\mathcal{L}_0$ with corners $(-L/2, -L)$, $(L/2, -L)$, $(L/2, L)$, $(-L/2, L)$. More precisely we define the field $\mathcal{Q}$ as:

$$\mathcal{Q}(r) = \begin{cases} \varphi^{(1)}(r) & \text{for } r \in \mathcal{L} \cap \mathcal{L}_0, \\ \varphi^{(2)}(-r) & \text{for } r \in (-\mathcal{L}) \cap \mathcal{L}_0. \end{cases} \quad (18)$$

We can implement now a discrete version of the m -reduction procedure ($m = 2$) by defining the fields:

$$\mathcal{X}(r) = \frac{1}{2} (\mathcal{Q}(r) + \mathcal{Q}(-r)), \quad (19)$$

$$\Phi(r) = \frac{1}{2} (\mathcal{Q}(r) - \mathcal{Q}(-r)), \quad (20)$$

where $r \in \mathcal{L}_0$. The resemblance with the continuum version of the m -reduction procedure is evident and the fields $\mathcal{X}$ and Φ are symmetric and antisymmetric with respect to the action of the generator $g : r \rightarrow -r$ of the discrete group Z_2 . The Hamiltonian in Eq. (4) can be rewritten in terms of these fields and, for $h = 0$, it becomes:

$$H = 2A \sum_{\langle r, r' \rangle \in \mathcal{L}} \cos(\mathcal{X}(r) - \mathcal{X}(r')) \cos(\Phi(r) - \Phi(r')), \quad (21)$$

which in the continuum limit corresponds to the action of our TM model:

$$\mathcal{A} = \int \left[\frac{1}{2} (\partial \mathcal{X})^2 + \frac{1}{2} (\partial \Phi)^2 \right] d^2x. \quad (22)$$

It is worth pointing out that the fields $\mathcal{X}$ and Φ are scalar fields and so the chiral fields defined by Eqs. (10), (11) can be seen as their chiral components. Moreover the group Z_2 is a discrete symmetry group, indeed both H and $\mathcal{A}$ are invariant under its action.

In the following we will show how the continuum version of the FFX model and its phase diagram can be described by the action:

$$\mathcal{A} = \int \left[\frac{1}{2} (\partial \mathcal{X})^2 + \frac{1}{2} (\partial \Phi)^2 + \mu \cos(\beta \Phi) + \lambda \cos\left(\frac{\beta}{2} \Phi + \delta\right) \right] d^2 x, \quad (23)$$

defined in terms of $m = 2$ reduced fields $\mathcal{X}$ and Φ , which embodies in its “neutral” sector the higher harmonic potential term conjectured in Refs. [1, 35, 41]. We assume the constraints $\beta^2 < 8\pi$, which characterizes both the cosine terms as relevant scalar fields perturbing the gaussian fixed point, and $|\delta| \leq \pi/2$ (see [44]). Thus the “neutral” sector is a two-frequency sine-Gordon theory that can be viewed as a deformation of a pure sine-Gordon one with the perturbing term $\lambda \cos(\beta \Phi/2 + \delta)$. The ultraviolet (UV) fixed point $\mu = 0, \lambda = 0$ of the action (23) corresponds to the TM model with central charge $c = 2$, describing the fixed point T in the phase diagram of the FFX model. Here the argument described in [44] can be adapted to our particular case to study the RG flow in the “neutral” sector of our theory. Let us define the dimensionless variable $\eta \equiv \lambda \mu^{-(8\pi - (\beta/2)^2)/(8\pi - \beta^2)}$; when $\eta = 0$ ($\lambda = 0$) the “neutral” sector reduces to a sine-Gordon model with a particle spectrum which consists of solitons and antisolitons and, for $\beta^2 < 4\pi$, some breathers. Switching on the perturbation (i.e. for $\lambda \neq 0$) a confinement of solitons into states with zero topological charge takes place and packets formed by 2 of the original solitons or antisolitons survive as stable excitations for generic values of $|\delta| < \pi/2$. In the limit $\eta \rightarrow \infty$ the 2-soliton evolves into the 1-soliton of the pure sine-Gordon model with $\mu = 0$. An unbinding phenomenon takes place in the particular $\delta = \pm\pi/2$ case for finite η and the 2-soliton decomposes into a sequence of two kinks K_1 . In such a case the limit $\eta \rightarrow \infty$ implies a transmutation of a composite topological excitation (the two kinks K_1) into an elementary one (the 1-soliton). So the existence of an intermediate critical value $\eta = \eta_c$ is required at which a phase transition takes place and the RG flow ends into the infrared (IR) fixed point described by a CFT with central charge $c = 1/2$, the Ising model. The central charge of the full model (23) so changes from $c = 2$ of the UV fixed point to $c = 3/2$ of the IR fixed point, i.e. we recover early known results of Monte Carlo simulations [28]. Indeed in the following we clarify how such an IR fixed point coincides with the $U(1) \otimes Z_2$ symmetric component ($c = 3/2$) of the whole TM model, which results then to properly describe the FFX model conjectured in Ref. [1], i.e. the fixed point P in the phase diagram of the FFX model. Then the full phase diagram of the FFX model in Fig. 1 would be recovered.

To such an extent we define fermion fields on two parallel layers in terms of the “neutral” boson fields. In particular let $\psi_i(x)$ be the fermion on layer i , whose chiral components at the UV fixed point are defined by:

$$(\psi_i)_{L/R} = \frac{1}{\sqrt{z}} : e^{i\alpha(\Phi)_{L/R}((-)^{i+1}z)} : ; \quad (24)$$

where $(\Phi)_{L/R}$ are the left and the right components of Φ . In terms of them the Z_2 symmetric ψ and Z_2 antisymmetric $\bar{\psi}$ fermions read as:

$$\psi = \frac{1}{2}(\psi_1 + \psi_2) \quad , \quad \bar{\psi} = \frac{1}{2}(\psi_1 - \psi_2). \quad (25)$$

The action (23) at $|\delta| = \pi/2$ can then be rephrased in terms of two interacting Ising layers as:

$$\mathcal{A} = \int \frac{1}{2} (\partial \mathcal{X})^2 d^2x + \mathcal{A}_1^{Ising} + \mathcal{A}_2^{Ising} + \rho \int d^2x (\varepsilon_1(x) \varepsilon_2(x)) + \quad (26)$$

$$+ \mu \int d^2x (\varepsilon_1(x) + \varepsilon_2(x)) + \lambda \int d^2x (\sigma_1(x) \sigma_2(x)), \quad (27)$$

where $\mathcal{A}_i^{Ising}$ denotes the fixed point action of the Majorana fermion ψ_i and $\varepsilon_i(x), \sigma_i(x)$ are the energy and spin of the i -th Ising model ($i = 1, 2$). This correspondence comes out observing that the two critical Ising models plus the interaction term in the coupling ρ define a line of $c = 1$ CFTs parameterized by ρ since the operator $\varepsilon_1 \varepsilon_2$ is marginal ($\Delta_{\varepsilon_1 \varepsilon_2} = 1$). In this one-parameter family of $c = 1$ CFTs the thermal $\varepsilon_1 + \varepsilon_2$ and the magnetic $\sigma_1 \sigma_2$ operators are known [45] to have ρ -dependent conformal dimensions which however satisfy the relation:

$$\frac{\Delta_{\sigma_1 \sigma_2}(\rho)}{\Delta_{\varepsilon_1 + \varepsilon_2}(\rho)} = \frac{\Delta_{\sigma_1 \sigma_2}(0)}{\Delta_{\varepsilon_1 + \varepsilon_2}(0)} = \frac{1}{4}. \quad (28)$$

It allows us to say that the thermal and magnetic terms in (27) play the role of the two cosine terms in (23), in particular the equivalence works under the following identifications:

$$\varepsilon_1 + \varepsilon_2 \sim \cos(\beta \Phi), \quad \sigma_1 \sigma_2 \sim \sin\left(\frac{\beta}{2} \Phi\right), \quad (29)$$

where the parameters ρ and β are linked by the relation $\Delta_{\varepsilon_1 + \varepsilon_2}(\rho) = \beta^2/8\pi$. Finally we are in the position to identify unambiguously the IR fixed point $c = 3/2$ of our theory. Indeed for each value of $\rho(\beta)$ there is a critical line in the $\mu\lambda$ -plane corresponding to the RG flow of our theory from the UV fixed point $c = 2$ to the IR fixed point $c = 3/2$; in particular we can consider the RG flow selected by the limit $\lambda \rightarrow \infty$. In such a case the magnetic term in (27) forces the spins σ_1 and σ_2 to align everywhere reducing the two Ising system to a single Ising one. Thus at the IR fixed point the degrees of freedom of the fermion $\bar{\psi}$ decouple and we are left with the $U(1) \otimes Z_2$ symmetric component ($c = 3/2$) of the whole TM model, i.e. the MR theory with partition function $Z^{MR}(w_c|\tau)$, as it can be immediately seen from Eq. (15).

5 Conclusions and outlooks

In this paper we proposed a $c = 2$ CFT description of the FFX model, which extends an early proposal by Foda [1]. A crucial role in such a construction is played by the Z_2

(Z_m in general) discrete symmetry built-in in the 2 (m in general)-reduction technique, which allows for non trivial topological properties of the vacuum.

By perturbing the $c = 2$ CFT with relevant operators, which provide higher harmonics contribution to the potential, it was shown how the unbinding of kink-antikink states can give rise to a massless line flow, which fully reproduces the phase diagram for the FFXY model. As a result of the flow the non invariant Z_2 degrees of freedom decouple, the partition function gets reduced to the Moore-Read one and the degeneracy of the ground state lost. The attractive infrared fixed point is identified with the $c = \frac{3}{2}$ superconformal field theory, that is a free massless field theory of one boson and one Majorana fermion, which well describes the critical behavior of the FFXY at the overlap temperature of the Ising and vortex unbinding transitions.

It is worth noticing that an interesting realization of the FFXY model physics just outlined can be given in terms of a two dimensional JJA in a transverse magnetic field [4, 14, 27, 38]. Such a system is a periodic array of superconducting islands connected by Josephson links; each island is characterized through the modulus and the phase of the order parameter. The lowering of the temperature to its critical value forces each island to become superconductive but the whole array is in a resistive state as long as the phases have not acquired a long range order. The array reaches such a global coherence state at a lower temperature and its behaviour can be well described only in terms of the order parameters phases φ_i through the Hamiltonian:

$$H_{JJA} = -E_J \sum_{\langle ij \rangle} \cos(\varphi_i - \varphi_j), \quad (30)$$

where $E_J > 0$ is the Josephson coupling energy and the sum is over nearest neighbours. Let us notice the similarity of Eq. (30) with the Hamiltonian of a two dimensional XY model. In this picture it is known that the array becomes superconducting below the temperature $T_J = \frac{\pi E_J}{2}$, where the phases are ordered and vortices may appear only in bounded pairs. Hence the ferromagnetic state for the XY model corresponds to the superconducting state for the JJA [13]. Such a regime is the classical one for a JJA, indeed the quantum fluctuations of the phases φ_i are weak and the vortices are well defined objects, which form a Coulomb gas and behave as particles with masses.

The analogy just outlined still works very well in the presence of a magnetic field perpendicular to the plane of the array, in this case a frustrated XY model can be realized; in particular, the full frustration $f = \frac{1}{2}$ corresponds to half a magnetic flux threading each cell. The ground state can be viewed as a pinned vortex lattice commensurate with the underlying periodic pinning potential leading to discrete symmetries, which add to the $U(1)$ symmetry of the superconducting order parameter [4, 14]. Furthermore we must remark the striking role of fully frustrated JJAs as an experimental tool for the study of phase transitions in FFXY type models [33, 34].

Acknowledgments. We would like to thank E. Orignac for his careful reading and comments on a preliminary version of this work. G. N. was supported by a postdoctoral fellowship of the Ministère français délégué à l'Enseignement supérieur et à la Recherche.

A TM on a torus

The partition function of the TM model on the torus has the following factorized form (see [3]):

$$Z(w_c|\tau) = Z^{MR}(w_c|\tau)Z_I(\tau), \quad (31)$$

in terms of the Moore-Read partition function Z^{MR} ($c = 3/2$) and of the Ising partition function Z_I ($c = 1/2$). They have the following expression:

$$Z^{MR}(w_c|\tau) = |\chi_0^{MC}(0|w_c|\tau)|^2 + |\chi_1^{MC}(0|w_c|\tau)|^2 + |\chi_2^{MC}(0|w_c|\tau)|^2, \quad (32)$$

$$Z_I(\tau) = |\bar{\chi}_0(\tau)|^2 + \left|\bar{\chi}_{\frac{1}{2}}(\tau)\right|^2 + \left|\bar{\chi}_{\frac{1}{16}}(\tau)\right|^2. \quad (33)$$

The above characters are defined by:

$$\bar{\chi}_0(\tau) = \frac{1}{2} \left(\sqrt{\frac{\theta_3(0|\tau)}{\eta(\tau)}} + \sqrt{\frac{\theta_4(0|\tau)}{\eta(\tau)}} \right), \quad (34)$$

$$\bar{\chi}_{\frac{1}{2}}(\tau) = \frac{1}{2} \left(\sqrt{\frac{\theta_3(0|\tau)}{\eta(\tau)}} - \sqrt{\frac{\theta_4(0|\tau)}{\eta(\tau)}} \right), \quad (35)$$

$$\bar{\chi}_{\frac{1}{16}}(\tau) = \sqrt{\frac{\theta_2(0|\tau)}{2\eta(\tau)}}, \quad (36)$$

which express the primary field content of the Ising model with Neveu-Schwartz (Z_2 twisted) boundary conditions, and by:

$$\chi_0^{MC}(0|w_c|\tau) = \chi_0(\tau)K_0(w_c|\tau) + \chi_{\frac{1}{2}}(\tau)K_2(w_c|\tau), \quad (37)$$

$$\chi_1^{MC}(0|w_c|\tau) = \chi_{\frac{1}{16}}(\tau) (K_1(w_c|\tau) + K_3(w_c|\tau)), \quad (38)$$

$$\chi_2^{MC}(0|w_c|\tau) = \chi_{\frac{1}{2}}(\tau)K_0(w_c|\tau) + \chi_0(0|\tau)K_2(w_c|\tau) \quad (39)$$

which express the primary field content of the Z_2 -invariant $c = \frac{3}{2}$ CFT. They are given in terms of a “charged” contribution ($c = 1$):

$$K_{2l+i}(w_c|\tau) = \frac{1}{\eta(\tau)} \Theta \left[\begin{array}{c} \frac{2l+i}{4} \\ 0 \end{array} \right] (2w_c|4\tau), \quad \forall (l, i) \in (0, 1)^2, \quad (40)$$

and a “isospin” one $\chi_\beta(\tau)$ ($c = 1/2$), where $w_c = \frac{1}{2\pi i} \ln z_c$ is the torus variable of the “charged” component.

References

- [1] O. Foda, *Nucl. Phys. B* **300** (1988) 611.
- [2] G. Cristofano, G. Maiella, V. Marotta, *Mod. Phys. Lett. A* **15** (2000) 1679.
- [3] G. Cristofano, G. Maiella, V. Marotta, G. Niccoli, *Nucl. Phys. B* **641** (2002) 547.
- [4] S. Teitel, C. Jayaprakash, *Phys. Rev. Lett.* **51** (1983) 1999.
- [5] T. C. Halsey, *Phys. Rev. B* **31** (1985) 5728.
- [6] B. I. Halperin, D. R. Nelson, *J. Low Temp. Phys.* **36** (1979) 1165.
- [7] T. C. Halsey, *J. Phys. C* **18** (1985) 2437; S. E. Korshunov, G. V. Uimin, *J. Stat. Phys.* **43** (1986) 1.
- [8] J. Villain, *J. Phys. C* **10** (1977) 1717.
- [9] G. Cristofano, G. Maiella, V. Marotta, *Mod. Phys. Lett. A* **15** (2000) 547.
- [10] G. Cristofano, V. Marotta, A. Naddeo, *Phys. Lett. B* **571** (2003) 250.
- [11] G. Cristofano, V. Marotta, A. Naddeo, *Nucl. Phys. B* **679** (2004) 621.
- [12] G. Cristofano, V. Marotta, A. Naddeo, *J. Stat. Mech.: Theor. Exp.* (2005) P03006; G. Cristofano, V. Marotta, A. Naddeo, G. Niccoli, *Eur. Phys. J. B* **49** (2006) 83.
- [13] J. M. Kosterlitz, D. J. Thouless, *J. Phys. C* **6** (1973) 1181; J. M. Kosterlitz, *J. Phys. C* **7** (1974) 1046.
- [14] S. Teitel, C. Jayaprakash, *Phys. Rev. B* **27** (1983) 598.
- [15] J. Villain, *J. de Physique* **36** (1975) 581; J. José, L. P. Kadanoff, S. Kirkpatrick, P. R. Nelson, *Phys. Rev. B* **16** (1977) 1217.
- [16] E. Fradkin, B. A. Huberman, S. H. Shenker, *Phys. Rev. B* **18** (1978) 4789.
- [17] S. Miyashita, H. Shiba, *J. Phys. Soc. Jpn.* **53** (1984) 1145; G. Grest, *Phys. Rev. B* **39** (1989) 9267; J. R. Lee, *Phys. Rev. B* **49** (1994) 3317; P. Olsson, *Phys. Rev. Lett.* **75** (1995) 2758; P. Olsson, *Phys. Rev. B* **55** (1997) 3585; S. Lee, K. C. Lee, *Phys. Rev. B* **49** (1994) 15184; Y. Ozeki, N. Ito, *Phys. Rev. B* **68** (2003) 054414.
- [18] M. Yosefin, E. Domany, *Phys. Rev. B* **32** (1985) 1778; J. M. Thijssen, H. J. F. Knops, *Phys. Rev. B* **42** (1990) 2438; J. Lee, J. M. Kosterlitz, E. Granato, *Phys. Rev. B* **43** (1991) R11531; E. Granato, M. P. Nightingale, *Phys. Rev. B* **48** (1993) 7438; G. Ramirez-Santiago, J. V. José, *Phys. Rev. Lett.* **68** (1992) 1224; E. H. Boubcheur, H. T. Diep, *Phys. Rev. B* **58** (1998) 5163.

- [19] Y. M. M. Knops, B. Nienhuis, H. J. F. Knops, H. W. J. Blote, *Phys. Rev. B* **50** (1994) 1061.
- [20] A. B. Zamolodchikov, V. A. Fateev, *Yad. Fiz.* **32** (1980) 581 [*Sov. J. Nucl. Phys.* **32** (1980) 298].
- [21] M. den Nijs, K. Rommelse, *Phys. Rev. B* **40** (1989) 4709.
- [22] M. P. Nightingale, H. W. J. Blote, *Phys. Rev. B* **33** (1986) 659; H. Schulz, T. Ziman, *Phys. Rev. B* **33** (1986) 6545; J. Solyom, T. Ziman, *Phys. Rev. B* **30** (1984) 3980.
- [23] U. Glaus, *Physica A* **141** (1987) 295; K. Saitoh, S. Takada, K. Kubo, *J. Phys. Soc. Jpn.* **56** (1987) 3755.
- [24] K. Nomura, *J. Phys. A* **28** (1995) 5451; A. Kitazawa, K. Nomura, K. Okamoto, *Phys. Rev. Lett.* **76** (1996) 4038; A. Kitazawa, K. Nomura, *J. Phys. Soc. Jpn.* **66** (1997) 3379; A. Kitazawa, K. Nomura, *J. Phys. Soc. Jpn.* **66** (1997) 3944; K. Nomura, A. Kitazawa, *J. Phys. A* **31** (1998) 7341.
- [25] W. Chen, K. Hida, B. C. Sanctuary, *Phys. Rev. B* **67** (2003) 104401.
- [26] S. E. Korshunov, *Phys. Rev. Lett.* **88** (2002) 167007.
- [27] M. Y. Choi, S. Doniach, *Phys. Rev. B* **31** (1985) 4516; E. Granato, J. M. Kosterlitz, J. Lee, M. P. Nightingale, *Phys. Rev. Lett.* **66** (1991) 1090; J. Lee, E. Granato, J. M. Kosterlitz, *Phys. Rev. B* **44** (1991) 4819; E. Granato, J. M. Kosterlitz, M. P. Nightingale, *Physica B* **222** (1996) 266.
- [28] M. P. Nightingale, H. W. J. Blote, *Phys. Rev. Lett.* **60** (1988) 1562; M. P. Nightingale, E. Granato, J. M. Kosterlitz, *Phys. Rev. B* **52** (1995) 7402.
- [29] P. Olsson, S. Teitel, *Phys. Rev. B* **71** (2005) 104423.
- [30] M. Hasenbusch, A. Pelissetto, E. Vicari, *J. Stat. Mech.: Theor. Exp.* (2005) P12002; M. Hasenbusch, A. Pelissetto, E. Vicari, *Phys. Rev. B* **72** (2005) 184502.
- [31] A. Jonsson, P. Minnhagen, *Phys. Rev. B* **55** (1997) 9035.
- [32] S. Bernhardsson, G. Cristofano, B. J. Kim, P. Minnhagen, work in preparation.
- [33] B. J. van Wees, H. S. J. van der Zant, J. E. Mooij, *Phys. Rev. B* **35** (1987) 7291.
- [34] C. J. Lobb, *Physica B* **126** (1984) 319; D. W. Abraham, C. J. Lobb, M. Tinkham, T. M. Klapwijk, *Phys. Rev. B* **26** (1982) 5268; R. F. Voss, R. A. Webb, *Phys. Rev. B* **25** (1982) 3446; C. J. Lobb, D. W. Abraham, M. Tinkham, *Phys. Rev. B* **27** (1983) 150.
- [35] K. Sokalski, T. Ruijgrok, B. Schoenmaker, *Physica A* **144** (1987) 322.

- [36] E. Granato, *J. Phys. C* **20** (1987) L215.
- [37] P. Baseilhac, *Nucl. Phys. B* **636** (2002) 465.
- [38] Y. M. Blanter, R. Fazio, G. Schon, *Nucl. Phys. B* **S58** (1997) 79.
- [39] E. Granato, J. M. Kosterlitz, *Phys. Rev. B* **33** (1986) 4767.
- [40] P. Minnhagen, *Phys. Rev. B* **32** (1985) 7548.
- [41] J. M. Thijssen, H. J. F. Knops, *Phys. Rev. B* **37** (1988) 7738.
- [42] L. B. Ioffe, V. B. Geshkenbein, M. V. Feigel'man, A. L. Fauchère, G. Blatter, *Nature* **398** (1999) 679; L. B. Ioffe, M. V. Feigel'man, *Phys. Rev. B* **66** (2002) 224503; B. Doucot, M. V. Feigel'man, L. B. Ioffe, *Phys. Rev. Lett.* **90** (2003) 107003; B. Doucot, L. B. Ioffe, J. Vidal, *Phys. Rev. B* **69** (2004) 214501.
- [43] X. G. Wen, Q. Niu, *Phys. Rev. B* **41** (1990) 9377; X. G. Wen, *Phys. Rev. B* **44** (1991) 2664; A. Yu. Kitaev, *Ann. Phys.* **303** (2003) 2.
- [44] G. Delfino, G. Mussardo, *Nucl. Phys. B* **516** (1998) 675.
- [45] L. P. Kadanoff, *Ann. Phys.* **120** (1979) 39; L. P. Kadanoff, A.C. Brown, *Ann. Phys.* **121** (1979) 318.